

引用文献

- 1) Sharma N, Cohen LG: Recovery of motor function after stroke. *Dev Psychobiol.* 2012; 54: 254-262.
- 2) Swayne OB, Rothwell JC, et al: Stages of motor output reorganization after hemispheric stroke suggested by longitudinal studies of cortical physiology. *Cereb Cortex*; 2008; 18: 1909-1922.
- 3) Blakemore SJ, Frith CD, et al: Spatio-temporal prediction modulates the perception of self-produced stimuli. *J Cogn Neurosci*; 1999; 11: 551-559.
- 4) Morioka S, Fukumoto T, et al: Changes in the equilibrium of standing on one leg at various life stages. *Curr Gerontol Geriatr Res.* 2012; 2012: 516283.
- 5) Morioka S, Yagi F: Influence of perceptual learning on standing posture balance: repeated training for hardness discrimination of foot sole. *Gait Posture*; 2004; 20: 36-40.
- 6) Morioka S, Yagi F: Effects of perceptual learning exercises on standing balance using a hardness discrimination task in hemiplegic patients following stroke: a randomized controlled pilot trial. *Clin Rehabil.*

2003;17: 600-607.

- 7) Morioka S, Hiyamizu M, et al: Effects of plantar hardness discrimination training on standing postural balance in the elderly: a randomized controlled trial. Clin Rehabil. 2009;23: 483-491.
- 8) Morioka S, Fujita H, et al: Effects of plantar perception training on standing posture balance in the old old and the very old living in nursing facilities: a randomized controlled trial. Clin Rehabil. 2011 ;25: 1011-1120.
- 9) Nakano H, Nozaki M, et al: Effect of a plantar perceptual learning task on walking stability in the elderly: a randomized controlled trial. Clin Rehabil. 2013; 27: 608-615.
- 10) Nakano H, Osumi M, et al: Changes in electroencephalographic activity during observation, preparation, and execution in a motor learning task. Int J Neurosci, 2014; 123: 866-75.
- 11) Wakata S, Morioka S: Brain activity and the perception of self-agency while viewing a video of tool manipulation: a functional near-infrared spectroscopy study. Neuroreport. 2014; 25: 422-426.

- 12)Wakata S, Morioka S: Brain activity and the perception of self-agency while viewing a video of hand grasping: a functional near-infrared spectroscopy study. *Neuroreport*. 2015; 26:394-398.
- 13)Hiyamizu M, Maeoka H, et al: Effects of video feedback on standing balance learning: A change of brain activity using functional near-infrared spectroscopy. *NeuroRehabilitation*. 2014; 35:579-585.
- 14)Hirakawa Y, Hara M, et al: Influence of psychological factors and neglect-like symptoms on postoperative pain after total knee arthroplasty. *Pain Res Manag*. 2014; 19:251-256.
- 15)Baliki MN, Geha PV, et al: A preliminary fMRI study of analgesic treatment in chronic back pain and knee osteoarthritis. *Mol Pain*. 2008; 4:47.
- 16)Apkarian AV, Sosa Y, et al: Chronic back pain is associated with decreased prefrontal and thalamic gray matter density. *J Neurosci*. 2004;24:10410-10415.
- 17)Maeoka H, Matsuo A, et al: Influence of transcranial direct current stimulation of the dorsolateral prefrontal cortex on pain related emotions:

- a study using electroencephalographic power spectrum analysis. *Neurosci Lett.* 2012; 512:12-16.
- 18) Ohmatsu S, Nakano H, et al: Activation of the serotonergic system by pedaling exercise changes anterior cingulate cortex activity and improves negative emotion. *Behav Brain Res.* 2014; 270:112-117.
- 19) Morioka S, Nobusako S, et al: Neurorehabilitation of chronic pain: Relationships among pain, motion, and perception. *Int J Phys Med Rehabil.* 2013; 1:132.
- 20) Vartiainen NV, Kirveskari E, et al. Central processing of tactile and nociceptive stimuli in complex regional pain syndrome. *Clin Neurophysiol.* 2008;119: 2380-2388.
- 21) Gieteling EW, van Rijn MA, et al: Cerebral activation during motor imagery in complex regional pain syndrome type 1 with dystonia. *Pain.* 2008; 134:302-329.
- 22) Moseley GL: I can't find it! Distorted body image and tactile dysfunction in patients with chronic back pain. *Pain.* 2008; 140:239-243.
- 23) Lewis JS, Schweinhardt P: Perceptions of the painful body: the

relationship between body perception disturbance, pain and tactile discrimination in complex regional pain syndrome. *Eur J Pain.* 2012;16:1320-1330.

24) Osumi M, Okuno H, et al: Tactile localization training for pain, sensory disturbance, and distorted body image: a case study of complex regional pain syndrome. *Neurocase.* 2015;21:628-634.

25) Katayama O, Iki H, et al: The effect of virtual visual feedback on supernumerary phantom limb pain in a patient with high cervical cord injury: a single-case design study. *Neurocase.* 2015: 1-7. [Epub ahead of print]

26) Nobusako S, Matsuo A, et al: Effectiveness of the Gaze Direction Recognition Task for Chronic Neck Pain and Cervical Range of Motion: A Randomized Controlled Pilot Study. *Rehabil Res Pract.* 2012; 2012:570387.

27) Moseley GL, Gallace A, et al: Bodily illusions in health and disease: physiological and clinical perspectives and the concept of a cortical 'body matrix'. *Neurosci Biobehav Rev.* 2012; 36: 34-46.

28) Osumi M, Imai R, et al: Factors associated with the modulation of pain by visual distortion of body size. *Front Hum Neurosci.* 2014; 8:137.

29) Osumi M, Imai R, et al: Negative body image associated with changes in the visual body appearance increases pain perception. *PLoS One.* 2014; 9:e107376.